arXiv:1510.00192v2 [math.CA] 4 Oct 2015

A note on a modified Bessel function integral

R. B. Paris
Division of Computing and Mathematics,
University of Abertay Dundee, Dundee DD1 1HG, UK

Abstract

We investigate the integral

$$\int_0^\infty \cosh^\mu t\, K_\nu(z\cosh t)\, dt \qquad \Re(z) > 0,$$

where K denotes the modified Bessel function, for non-negative integer values of the parameters μ and ν. When the integers are of different parity, closed-form expressions are obtained in terms of $z^{-1}e^{-z}$ multiplied by a polynomial in z^{-1} of degree dependent on the sign of $\mu-\nu$.

Mathematics Subject Classification: 30E20, 33C10

Keywords: Modified Bessel function, integrals, spherical Bessel function

1. Introduction

An integral arising in neutron scattering [1, 2] is given by

$$\mathcal{F}(\mu,\nu;z) := \int_0^\infty \cosh^\mu t\, K_\nu(z\cosh t)\, dt \qquad \Re(z) > 0, \tag{1.1}$$

where $K_\nu(z)$ is the modified Bessel function of the second kind with the parameters $\mu \geq 0$ and $\nu \in C$. The asymptotic expansion of $\mathcal{F}(\mu,\nu;z)$ for $|z| \to \infty$ in $\Re(z) > 0$ has been obtained by Birrell in [1]. The integral (1.1) can also be expressed in terms of a combination of three ${}_1F_2(z^2/4)$ generalised hypergeometric functions. The large-z asymptotics could therefore be derived by means of the asymptotic theory of integral functions of hypergeometric type; see, for example, the summary presented in [5, Section 2.3].

Birrell [1] placed particular emphasis on the special case when μ and ν are positive integers of different parity, where $\mathcal{F}(\mu,\nu;z)$ reduces to a finite expansion in inverse powers of z multiplied by $z^{-1}e^{-z}$. The approach adopted in [1] was the use of a recurrence relation satisfied by the integral in (1.1) combined with direct evaluation when $\mu = 0$, ν odd and $\mu = 1$, ν even. In this note, we consider the case of integer μ and ν from a different viewpoint.

2. Evaluation for non-negative integer μ

We consider the evaluation of $\mathcal{F}(\mu,\nu;z)$ when μ and ν are non-negative integers of different parity by using a well-known property of spherical Bessel functions.

2.1 *The case of even μ*

Let us first consider $\mu=2n$ and $\nu=2m+1$, where m and n are non-negative integers. Then we have [3, p. 136]

$$\cosh^{2n}t=\frac{1}{2^{2n-1}}\sum_{k=0}^{n}{}'\binom{2n}{k}\cosh 2(n-k)t,$$

where the prime after the summation sign indicates that the term $k=n$ is multiplied by the factor $\frac{1}{2}$. We also make use of the result [4, p. 253]

$$\int_0^\infty \cosh(a-b)t\,K_{a+b}(2x\cosh t)\,dt=\tfrac{1}{2}K_a(x)K_b(x)\qquad(\Re(x)>0;\ a,b\in C).$$

It therefore follows that

$$\mathcal{F}(2n,2m+1;z)=\frac{1}{2^{2n}}\sum_{k=0}^{n}{}'\binom{2n}{k}K_{m+n-k+\frac{1}{2}}(\tfrac{1}{2}z)K_{m-n+k+\frac{1}{2}}(\tfrac{1}{2}z),\tag{2.1}$$

where the Bessel functions are of semi-integer order and so have finite expressions given by [4, p. 264]

$$K_{s+\frac{1}{2}}(x)=\sqrt{\frac{\pi}{2x}}\,e^{-x}\sum_{k=0}^{s}\frac{(s+k)!}{k!(s-k)!}\,(2x)^{-k}\qquad(s=0,1,2,\ldots).$$

Then, when $m\geq n$ we obtain

$$\begin{aligned}\mathcal{F}(2n,2m+1;z)&=\frac{\pi e^{-z}}{2^{2n}z}\sum_{k=0}^{n}{}'\binom{2n}{k}\sum_{r=0}^{m+n-k}a_r z^{-r}\sum_{s=0}^{m-n+k}b_s z^{-s}\qquad(2.2)\\&=\sum_{k=0}^{n}{}'\binom{2n}{k}\sum_{p=0}^{2m}c_p(k)z^{-p},\end{aligned}$$

where

$$a_r=\frac{(m+n-k+r)!}{r!(m+n-k-r)!},\qquad b_s=\frac{(m-n+k+s)!}{s!(m-n+k-s)!}\tag{2.3}$$

and we have defined

$$\begin{aligned}c_p(k)&\equiv c_p(k;m,n)=\sum_{r+s=p}a_r b_s\\&=\sum_{r=0}^{p}\frac{(m+n-k+r)!(m-n+k+p-r)!}{r!(p-r)!(m+n-k-r)!(m-n+k-p+r)!}.\end{aligned}$$

We therefore find the result

$$\mathcal{F}(2n,2m+1;z)=\frac{\pi e^{-z}}{2^{2n}z}\sum_{p=0}^{2m}C_p(m,n)z^{-p}\qquad(m\geq n),\tag{2.4}$$

where the coefficients $C_p(m,n)$ are given by

$$C_p(m,n) = \sum_{k=0}^{n}{}' \binom{2n}{k} c_p(k) \qquad (0 \le p \le 2m). \tag{2.5}$$

In the case $m < n$, we can proceed in a similar manner but now care has to be taken with the second Bessel function appearing in (2.1). Making use of the fact that $K_{-\nu}(x) = K_\nu(x)$, we write

$$K_{m-n+k+\frac{1}{2}}(\tfrac{1}{2}z) = \begin{cases} K_{n-m-k-\frac{1}{2}}(\frac{1}{2}z) & \text{for } k \le n-m-1 \\ \\ K_{m-n+k+\frac{1}{2}}(\frac{1}{2}z) & \text{for } k \ge n-m. \end{cases} \tag{2.6}$$

Then the product in (2.2) becomes

$$\sum_{r=0}^{m+n-k} a_r z^{-r} \sum_{s=0}^{S} b_s z^{-s}, \qquad S = \begin{cases} n-m-k-1 & \text{for } k \le n-m-1 \\ m-n+k & \text{for } k \ge n-m \end{cases}$$

to yield

$$\mathcal{F}(2n, 2m+1; z) = \frac{\pi e^{-z}}{2^{2n} z} \sum_{k=0}^{n}{}' \binom{2n}{k} \sum_{p=0}^{2n-1} d_p(k) z^{-p}.$$

The coefficients $d_p(k) \equiv d_p(k; m, n)$ are defined by

$$d_p(k) = \sum_{r+s=p} a_r \hat{b}_s, \qquad \hat{b}_s = \begin{cases} \dfrac{(n-m-k-1+s)!}{s!(n-m-k-1-s)!} & \text{for } k \le n-m-1 \\ \\ \dfrac{(m-n+k+s)!}{s!(m-n+k-s)!} & \text{for } k \ge n-m, \end{cases} \tag{2.7}$$

where the a_r are as specified in (2.3).

Consequently, we obtain the result

$$\mathcal{F}(2n, 2m+1; z) = \frac{\pi e^{-z}}{2^{2n} z} \sum_{p=0}^{2n-1} D_p(m,n) z^{-p} \qquad (m < n), \tag{2.8}$$

where the coefficients $D_p(m,n)$ are given by

$$D_p(m,n) = \sum_{k=0}^{n}{}' \binom{2n}{k} d_p(k) \qquad (0 \le p \le 2n-1). \tag{2.9}$$

2.2 The case of odd μ

When $\mu = 2n+1$, we can make use of the expansion [3, p. 136]

$$\cosh^{2n+1} t = \frac{1}{2^{2n}} \sum_{k=0}^{n} \binom{2n+1}{k} \cosh(2n-2k+1)t$$

to obtain

$$\mathcal{F}(2n+1, 2m; z) = \frac{1}{2^{2n+1}} \sum_{k=0}^{n} \binom{2n+1}{k} K_{m+n-k+\frac{1}{2}}(\tfrac{1}{2}z) K_{m-n+k-\frac{1}{2}}(\tfrac{1}{2}z). \tag{2.10}$$

Following the same procedure as in the even μ case, we obtain when $m > n$

$$\mathcal{F}(2n+1,2m;z) = \frac{\pi e^{-z}}{2^{2n+1}z}\sum_{k=0}^{n}\binom{2n+1}{k}\sum_{p=0}^{2m-1}\hat{C}_p(m,n)z^{-p} \qquad (m>n), \tag{2.11}$$

where

$$\hat{C}_p(m,n) = \sum_{k=0}^{n}\binom{2n+1}{k}\hat{c}_p(k) \qquad (0\le p\le 2m-1), \tag{2.12}$$

with

$$\hat{c}_p(k) = \sum_{r+s=p}\frac{(m+n-k+r)!(m-n+k-1+s)!}{r!s!(m+n-k-r)!(m-n+k-1-s)!}.$$

When $m \le n$, the second Bessel function in (2.10) is written as in (2.14) with k replaced by $k-1$ to find

$$\mathcal{F}(2n+1,2m;z) = \frac{\pi e^{-z}}{2^{2n+1}z}\sum_{k=0}^{n}\binom{2n+1}{k}\sum_{p=0}^{2n}\hat{D}_p(m,n)z^{-p} \qquad (m\le n), \tag{2.13}$$

where

$$\hat{D}_p(m,n) = \sum_{k=0}^{n}\binom{2n+1}{k}\hat{d}_p(k) \qquad (0\le p\le 2n), \tag{2.14}$$

with $\hat{d}_p(k)$ as defined in (2.7) with $k\to k-1$.

We summarise the results obtained in the following theorem.

Theorem 1. *Let m and n be non-negative integers and $\Re(z)>0$. Then*

$$\mathcal{F}(2n,2m+1;z) = \frac{\pi e^{-z}}{2^{2n}z}\,P(z), \qquad \mathcal{F}(2n+1,2m;z) = \frac{\pi e^{-z}}{2^{2n+1}z}\,\hat{P}(z), \tag{2.15}$$

where $P(z)$ and $\hat{P}(z)$ are polynomials in z^{-1} given by

$$P(z) = \begin{cases} \displaystyle\sum_{p=0}^{2m} C_p(m,n)z^{-p} & (m\ge n) \\ \displaystyle\sum_{p=0}^{2n-1} D_p(m,n)z^{-p} & (m<n) \end{cases}, \qquad \hat{P}(z) = \begin{cases} \displaystyle\sum_{p=0}^{2m} \hat{C}_p(m,n)z^{-p} & (m>n) \\ \displaystyle\sum_{p=0}^{2n} \hat{D}_p(m,n)z^{-p} & (m\le n). \end{cases}$$

The coefficients $C_p(m,n)$, $D_p(m,n)$, $\hat{C}_p(m,n)$ and $\hat{D}_p(m,n)$ are defined in (2.5), (2.9), (2.12) and (2.14).

3. Concluding remarks

As an example, when $\mu=4$, $\nu=7$ ($m=3$, $n=2$) and $\mu=4$, $\nu=3$ ($m=1$, $n=2$) the values of the coefficients $C_p(m,n)$ are

$$8,\quad 208,\quad 2520,\quad 17880,\quad 76800,\quad 184320,\quad 184320 \qquad (m=3,n=2;\ \ 0\le p\le 2m),$$

$$8,\quad 48,\quad 120,\quad 120 \qquad (m=1,n=2;\ \ 0\le p\le 2n-1);$$

whence we find

$$\mathcal{F}(4,7;z) = \frac{\pi e^{-z}}{2z}\,(1 + 26z^{-1} + 315z^{-2} + 2235z^{-3} + 9600z^{-4} + 23040z^{-5} + 23040z^{-6})$$

and

$$\mathcal{F}(4,3;z) = \frac{\pi e^{-z}}{2z}\,(1 + 6z^{-1} + 15z^{-2} + 15z^{-3}).$$

We note from Theorem 1 that the finite series in z^{-1} terminate at different p index. For even μ, the degree of the polynomial $P(z)$ is $2m$ when $m \geq n$ and $2n-1$ when $m < n$. For odd μ, the degree of the polynomial $\hat{P}(z)$ is $2m$ when $m > n$ and $2n$ when $m \leq n$. This was observed in [1], although the case $m = n$ was not mentioned.

Since it is easily verified that $c_0(k) = d_0(k) = \hat{c}_0(k) = \hat{d}_0(k) = 1$, the leading coefficients in the expansions in (2.15) satisfy

$$C_0(m,n) = D_0(m,n) = \sum_{k=0}^{n}{}'\binom{2n}{k} = 2^{2n} - \frac{n!}{2(\frac{1}{2})_n}\binom{2n}{n},$$

$$\hat{C}_0(m,n) = \hat{D}_0(m,n) = \sum_{k=0}^{n}\binom{2n+1}{k} = 2^{2n},$$

which are independent of m.

Finally, we mention that the integral

$$\mathcal{G}(2n,\nu;z) = \int_0^\infty \sinh^{2n}\!t\, K_\nu(z\cosh t)\,dt$$

when $\nu = 2m+1$ can be dealt with in the same manner. Since [3, p. 136]

$$\sinh^{2n}\!t = \frac{1}{2^{2n-1}}\sum_{k=0}^{n}{}'(-)^k\binom{2n}{k}\cosh 2(n-k)t,$$

it follows immediately from Section 2.1 that

$$\mathcal{G}(2n,2m+1;z) = \frac{\pi e^{-z}}{2^{2n}z}\,P(z),$$

where the polynomial $P(z)$ is as specified in Theorem 1, with the coefficients $C_p(m,n)$ and $D_p(m,n)$ now defined by

$$C_p(m,n) = \sum_{k=0}^{n}{}'(-)^k\binom{2n}{k}c_p(k), \qquad D_p(m,n) = \sum_{k=0}^{n}{}'(-)^k\binom{2n}{k}d_p(k).$$

The leading coefficients are again independent of m and given by

$$C_0(m,n) = D_0(m,n) = \sum_{k=0}^{n}{}'(-)^k\binom{2n}{k} = 0;$$

this means that a z^{-1} can be factored out leaving the polynomial $P(z)$ with degree reduced by unity.

References

[1] J. Birrell, Evaluation of a family of Bessel function integrals. (2015) arXiv:1509.06308.

[2] J. Birrell, C.T. Yang and J. Rafelski, Relic neutrino freeze-out: dependence on natural constants. Nucl. Phys. B **890** (2015) 481.

[3] A. Jeffrey and H-H. Dai, *Handbook of Mathematical Formulas and Integrals*, 4th edition, Academic Press, Oxford, 2008.

[4] F.W.J. Olver, D.W. Lozier, R.F. Boisvert and C.W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.

[5] R.B. Paris and D. Kaminski, *Asymptotics and Mellin-Barnes Integrals*, Cambridge University Press, Cambridge, 2001.